



## A QUANTUM MECHANICAL STUDY ON CH<sub>2</sub>NS<sup>+</sup> FAMILY OF CATIONS, POSSIBLE INTERSTELLAR SPECIES

M. Gronowski<sup>1</sup> and R. Kołos<sup>1,2</sup>

**Abstract.** It is demonstrated on quantum mechanical grounds that the dissociative recombination of HSCNH<sup>+</sup> with an electron, a reaction of interest to the chemistry of molecular interstellar clouds, leads either to HSCN or to HNCS. The HSCNH<sup>+</sup> cation, thus far undiscovered in space, is therefore proposed as a precursor for both neutral interstellar species, while H<sub>2</sub>NCS<sup>+</sup> (32 kJ/mol more stable than HSCNH<sup>+</sup>, according to our quantum-chemical calculations) can yield only HNCS. Calculations of potential energy surfaces were carried out for both cationic precursors.

### 1 Introduction

HNCS and HSCN were predicted as the lowest energy structures among the molecules sharing the same stoichiometry, with HSCN being 26 kJ/mol less stable than HNCS (Wierzejewska & Moc 2003). The importance of HSCN would therefore be negligible if thermodynamic equilibrium between the two species existed. Both molecules are nevertheless detected in the interstellar medium (Frerking *et al.* 1979; Halfen *et al.* 2009), and the HSCN abundance in the prominent molecular cloud Sgr B2 is only three times lower than that of HNCS (Adande *et al.* 2010; Halfen *et al.* 2009). Adande *et al.* (2010) proposed cationic precursors for the following reactions:



The identification of H<sub>2</sub>NCS<sup>+</sup> or its isomers is crucial for uncovering the HNCS/HSCN astrochemistry.

---

<sup>1</sup> Institute of Physical Chemistry, Polish Academy of Sciences, 44/52 Kasprzaka, 01-224 Warsaw, Poland; e-mail: [marcingronowski@gmail.com](mailto:marcingronowski@gmail.com); [rkolos@ichf.edu.pl](mailto:rkolos@ichf.edu.pl)

<sup>2</sup> Cardinal S. Wyszyński University, Faculty of Mathematics and Natural Sciences, Wóycickiego 1/3, 01-938 Warsaw, Poland; e-mail: [rkols@ichf.edu.pl](mailto:rkols@ichf.edu.pl)

Here we report the first results of a theoretical study on relevant dissociative recombination processes for  $\text{H}_2\text{NCS}^+$  isomers.

## 2 Theoretical methods




The Gaussian 03 (Frisch *et al.* 2004) software package was used throughout this study. Preliminary structures of  $\text{H}_2\text{NCS}^+$  isomers were indicated by the density functional theory, B3LYP/aug-cc-pVTZ (Becke 1993; Kendall *et al.* 1992; Lee *et al.* 1988). The coupled cluster theory level (Kendall *et al.* 1992; Pople *et al.* 1987) with single and double (CCSD/cc-pVTZ) excitations was employed to perform the final geometry optimizations and the potential energy surface (PES) scans.


## 3 Results and discussion

Structures corresponding to the deepest minima on the singlet potential energy surface of  $\text{CH}_2\text{NS}^+$  are presented in Figure 1. Vibrational frequency calculations allowed for the distinction between true minima and saddle points. The radio astronomical detection of the most stable singlet isomers **1** and **2** seems to be feasible, given their predicted electric dipole moments of 3 D.


Within a quiescent interstellar cloud, dissociative recombination consists of the collision of a cation  $\text{M}^+$  with a slow electron, which creates M in one of many highly excited electronic states, near the ionisation limit. Neutral M is then subjected to a range of different unimolecular transformations governed by potential energy surfaces of its ground or excited states. The simplification applied here, as well as in similar studies (Kolos *et al.* 2009; Osamura *et al.* 1999), consists in reducing the problem to the ground PES. While the neglect of rearrangements taking place in excited electronic states removes severe computational difficulties, it has to be regarded as a preliminary approach, showing the fate of only those M species which would rapidly pass to the ground state. The coupling between excited and ground states is in fact not expected to be strong for a 5-atom molecule. We have analysed, along these lines, the dissociation of the radical species  $\text{H}_2\text{NCS}$  and  $\text{HNCSH}$  (denoted **R1** and **R2** in Fig. 2).

Geometries of flat **R1** and **R2** radicals are significantly different from those of their respective cations **1** and **2**, yet the energetic separations **R1-R2** and **1-2** are very similar. The CCSD/cc-pVTZ energy of **R2** (ZPE-corrected) is 40 kJ/mol higher than that of **R1**. The ionisation energy of both radicals is approx. 645 kJ/mol. Radicals **R1** and **R2** may yield several species, namely:





**Fig. 1.** CH<sub>2</sub>NS<sup>+</sup> cationic structures corresponding to the lowest energy minima of the singlet PES. Isomer labels and symmetry designations are followed by CCSD/cc-pVTZ relative energies (kJ/mol; corrected for the zero-point vibrational energy, calculated with respect to the most stable species; 1 eV  $\approx$  96 kJ/mol), and by electric dipole moment values.



**Fig. 2.** Geometries of H<sub>2</sub>NCS and HNC<sub>2</sub>SH radicals (Å, degrees) calculated by CCSD/cc-pVTZ.



The activation energy  $E_a$  is predicted to be the lowest for the dissociation of radical **R2** into HNC and SH. This reaction, while important as an **R2** destruction route, cannot significantly increase the amount of abundant interstellar HNC or SH molecules. Of note, **R2** may yield HNCS and HSCN. It can also, formally, dissociate into HN and CSH, but this latter reaction is not likely, with its predicted  $E_A$  value of 684 kJ/mol, *i.e.* higher than the excess energy with which the neutral radical is produced.

In the case of **R1**, the lowest activation energy is predicted from the dissociation of **R1** into HNCS and H. The  $E_a$  values are significantly higher for other processes. The calculated heights of energetic barriers indicate that HNCS should be produced more efficiently than HSCN, in accordance with interstellar abundance measurements.

## 4 Summary

HNCS and NCSH can be formed in the interstellar medium from the two cationic precursors:  $\text{HNCSH}^+$  and  $\text{H}_2\text{NCS}^+$ . The calculated heights of the energy barriers indicate that HNCS should be produced more efficiently than HSCN, in qualitative consistence with interstellar abundance measurements. Molecules HNC, HS,  $\text{H}_2\text{NC}$ , CS and  $\text{NH}_2$  can arise in associated dissociative recombinations. Y-shaped  $\text{H}_2\text{NCS}^+$  and L-shaped  $\text{HNCSH}^+$  species are the most stable isomers of this stoichiometry, and, with their predicted electric dipole moments of 3 D, are likely to be discovered with microwave spectroscopy.

This work was supported by the Polish Ministry of Science and Education, project No. N N204 152 036. Quantum chemical calculations were accomplished at the Interdisciplinary Centre for Mathematical and Computational Modelling (grant G36-13). M.G. acknowledges a *START* stipend granted by The Foundation for Polish Science.

## References

- Adande, G.R., Halfen, D.T., Ziurys, L.M., Quan, D., & Herbst, E., 2010, *ApJ*, 725, 561  
Becke, A.D., 1993, *J. Chem. Phys.*, 98, 5648  
Frerking, M.A., Linke, R.A., & Thaddeus, P., 1979, *ApJ*, 234, L143  
Frisch, M.J., Trucks, G.W., Schlegel, H.B., *et al.*, 2004, Gaussian 03, Gaussian, Inc., Wallingford, CT  
Halfen, D.T., Ziurys, L.M., Brunken, S., *et al.*, 2009, *ApJ*, 702, L124  
Kendall, R.A., Dunning, T.H., & Harrison, R.J., 1992, *J. Chem. Phys.*, 96, 6796  
Kolos, R., Gronowski, M., & Dobrowolski, J.C., 2009, *ApJ*, 701, 488  
Lee, C., Yang, W., & Parr, R.G., 1988, *Phys. Rev. B*, 37, 785  
Osamura, Y., Fukuzawa, K., Terzieva, R., & Herbst, E., 1999, *ApJ*, 519, 697  
Pople, J.A., Head-Gordon, M., & Raghavachari, K., 1987, *J. Chem. Phys.*, 87, 5968  
Wierzejewska, M., & Moc, J., 2003, *J. Phys. Chem.*, 107, 11209